本種ハこちくノ類ニシテこちくトハ薬鞘ニ逆毛ナキ事薬裏ニハ唯半面ニ散生 スル毛アルノミ故區別サレ、よもぎだこちくヨリハ節ニ細カナル密毛アルト薬 質カタキ事、薬脚ハ雨縁ニ鬢毛アル事トニテ區別シ得、信濃國下伊那郡豐村村 ニテ杉本順一氏發見採集ス。

## **E**RRATA

- p. 568, between line 28 and 29 put 'Arundinaria viridistriata var. vagans Nakai, comb. nov'., and remove line 29 under line 33 as a synonym, changing gothics to italics.
- p. 569, line 9 for Arundinaria Hisauchii Nakai read Arundinaria hakonensis Nakai, nom. nov.
- p. 570, line 17. for Uchiyama read Uchida.

## Studies on the Lichens of Japan (IV)

Ву

M. M. Satô

佐藤正己: 日本産地衣類ノ研究 (其四)

## Notes on Usnea diffracta WAINIO

Usnea diffracta Wainio in Bot. Mag. Tokyo XXXV, p. 45 (1921). Motyka in Zahlbruckner in Fedde Repertorium XXXIII, p. 62 (1933). Syn. Usnea plicata var. annulata Müll. Arg. in Nuov. Giorn. Bot. Ital. XXIV, p. 191 (1892). Asahina in Journ. Japan. Bot. VIII, p. 252, (1929)—syn. nov.

Usnea dasypoga var. plicata f. annulata Müll. Arg. apud Hue, Lich. Extra-Europ., no. 126, T. I, Pl. III, fig. 6 (1901)—syn. nov.

Usnea longissima (non ACH.) MIYOSHI, Cryptog. Japon. Icon. Illust., I, Pl. IV (1899).

In describing *Usnea diffracta* as a new species, Wainio might have overlooked the fact that the same lichen was described already as a variety